

## Dog Vaccination SCHEDULE

AGE	VACCINATION
<b>5 wks</b>	<b>Parvovirus:</b> For puppies at high risk. Check with your local veterinarian.
<b>6 &amp; 9 wks</b>	<b>Combination vaccine*</b> without leptospirosis. <b>Coronavirus:</b> Where coronavirus is a concern.
<b>12 wks or older</b>	<b>Rabies:</b> Given by your local veterinarian (age at vaccination may vary according to local law).
<b>12 &amp; 15 wks</b>	<b>Combination vaccine*:</b> Include leptospirosis where it is a concern, or if traveling to an area where it occurs. <b>Coronavirus:</b> Where coronavirus is a concern. <b>Lyme:</b> Where Lyme disease is a concern or if traveling
<b>Adult</b>	<b>Combination vaccine*:</b> Include leptospirosis where it is a concern, or if traveling to an area where it occurs. <b>Coronavirus:</b> Where coronavirus is a concern. <b>Lyme:</b> Where Lyme disease is a concern or if traveling to an area where it occurs. <b>Rabies:</b> Given by your local veterinarian (time interval between vaccinations may vary according to local law).

These recommendations vary depending on the age, breed, and health status of the dog, the potential of the dog to be exposed to the disease, the type of vaccine, whether the dog is used for breeding, and the geographical area where the dog lives or may visit.

\*A combination vaccine, often called a 5- way vaccine, usually includes adenovirus cough and hepatitis, canine distemper, parainfluenza, and parvovirus. Some combination vaccines may also include leptospirosis (7-way vaccines) and/or coronavirus. The inclusion of either canine adenovirus-1 or adenovirus- 2 in a vaccine will protect against both adenovirus cough and hepatitis; adenovirus-2 is highly preferred.

\*\*Some puppies may need additional vaccinations against parvovirus after 15 weeks of age. Consult with your local veterinarian. Bordetella and parainfluenza: For complete canine cough protection, we recommend Intra-Trac III ADT . For dogs that are shown, in field trials, or are boarded, we recommend vaccination every six to twelve months with Intra-Trac III ADT .

### Use needles & syringes only once!

With any vaccine or injectable medication, always use a separate sterile needle and syringe for each injection. Safely dispose of all used syringes and needles.